



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE FUNCTIONAL EQUATION FOR THE SINE. ADDITIONAL NOTE.

BY EDWARD B. VAN VLECK.

In vol. 10 of the Annals, p. 161, I brought forward the following functional equation for the sine:

$$(I) \quad f(x - y + A) - f(x + y + A) = 2f(x)f(y).$$

In the second of the two proofs there given for the definition of the sine by means of this equation, one step should have been supplied to the reader between 7) and 8) to justify in the second equation of 8) the exclusion of the values $f(x) = \pm 1$ for $0 < x < B < A$. The justification of this exclusion runs as follows.

Suppose k to be a value between 0 and A for which $f(k) = \pm 1$. Then by my equation * (2),

$$f(k + A) = 0.$$

Putting $y = k$ in (1) we have

$$f(x - k + A) + f(x + k + A) = 0.$$

But for $y = k$ equation (I) becomes

$$f(x - k + A) - f(x + k + A) = \pm 2f(x).$$

The combination of these two equations gives

$$f(x + k + A) = \mp f(x).$$

This is equivalent to

$$f(x + k - A) = \pm f(x),$$

since the addition of $2A$ to the argument of the function changes its sign. Hence if x is replaced in (I) by $x + k - A$, there results the equation

$$f(x - y + k) - f(x + y + k) = \pm 2f(x)f(y),$$

where the upper or lower sign holds in the right hand member according as $f(x) = +1$ or $f(x) = -1$. In the former case the equation is the same as (I) with k in place of A , while in the latter case we have merely to put $f_1(x) \equiv -f(x)$ to make it the same. We may therefore suppose A in (I) to be the smallest value of $k > 0$ for which $f(k) = \pm 1$. In other words, between 0 and A we have

$$-1 < f(x) < 1.$$

* The equations referred to are

$$(1) \quad f(x - y + A) + f(x + y + A) = 2f(x + A)f(y + A),$$

$$(2) \quad 1 = f^2(x) + f^2(x + A).$$